

DIRECT AND INVERSE UNITARY EIGENPROBLEMS IN SIGNAL PROCESSING: AN OVERVIEW ¹

G.S. Ammar
Department of Mathematical Sciences
Northern Illinois University
DeKalb, IL 60115 USA
ammar@math.niu.edu

W.B. Gragg
Department of Mathematics
Naval Postgraduate School
Monterey, CA 93940 USA
gragg@guinness.math.nps.navy.mil

L. Reichel
Department of Mathematics and Computer Science
Kent State University
Kent, OH 44242 USA
reichel@mcs.kent.edu

KEYWORDS. Unitary Hessenberg matrices, Szegő polynomials, frequency estimation, updating and downdating least-squares approximants, eigenvalue computations, inverse eigenvalue problem.

Let \mathcal{H}_n denote the set of unitary upper Hessenberg matrices of order n with nonnegative subdiagonal elements. These matrices bear many similarities with real symmetric tridiagonal matrices, both in terms of their structure, their underlying connections with orthogonal polynomials, and the existence of efficient algorithms for solving eigenproblems for these matrices. The *Schur parameterization* of \mathcal{H}_n provides the means for the development of efficient algorithms for finding eigenvalues and eigenvectors of these matrices. These algorithms include the *QR* algorithm for unitary Hessenberg matrices [9], an algorithm for solving the orthogonal eigenproblem using two half-size singular value decompositions [1], a divide-and-conquer method [10, 5], an approach based on matrix pencils [6], and a unitary analog of the Sturm sequence method [7]. Aspects of inverse eigenproblems for unitary Hessenberg matrices are considered in [3] and efficient algorithms for constructing a unitary Hessenberg matrix from spectral data are presented in [13, 3].

Unitary Hessenberg matrices are fundamentally connected with Szegő polynomials; i.e., with polynomials orthogonal with respect to a measure on the unit circle in the complex plane [8]. In particular, the Schur parameters of a unitary Hessenberg matrix $H \in \mathcal{H}_n$

¹This paper appears in *Linear Algebra for Large Scale and Real-Time Applications*, M. S. Moonen, G. H. Golub and B. L. R. De Moor, eds., Kluwer Academic Publishers, Dordrecht, The Netherlands, 1993, pp. 341–343.

are the recurrence coefficients of the Szegő polynomials determined by a discrete measure on the unit circle. Moreover, the monic Szegő polynomial of degree k is the characteristic polynomial of the leading principal submatrix H_k of $H = H_n$. The zeros of the Szegő polynomial of degree n , which are the eigenvalues of H , are the nodes of the discrete measure. Furthermore, the weights of the measure are proportional to the squared moduli of the first components of the normalized eigenvectors of H . Algorithms for eigenproblems and inverse eigenproblems for unitary Hessenberg matrices are therefore useful in several computational problems in signal processing involving Szegő polynomials.

Eigenproblems for unitary Hessenberg matrices naturally arise in several frequency estimation procedures in signal processing, including Pisarenko's method [2] and the composite sinusoidal modeling method [11].

In [12], an efficient and reliable algorithm is presented for discrete least-squares approximation on the unit circle by algebraic polynomials. This algorithm also applies to the special case of discrete least-squares approximation of a real-valued function given at arbitrary distinct nodes in $[0, 2\pi)$ by trigonometric polynomials. The algorithm is based on the solution of an inverse eigenproblem for unitary Hessenberg matrices presented in [3]. This algorithm is an *updating* procedure in that the least-squares approximant is obtained by incorporating the nodes of the inner product one at a time. Numerical experiments show that the algorithm produces consistently accurate results that are often better than those obtained by general QR decomposition methods for the least-squares problem.

In [4] an algorithm is presented for *downdating* the Szegő polynomials and given least-squares approximant when a node is deleted from the inner product. This scheme uses the unitary Hessenberg QR algorithm [9] with the node to be deleted as a shift. This algorithm can be combined with the fast Fourier transform (FFT) when the given nodes are a subset of equispaced points. This situation arises, for example, when an FFT is performed on a data set with some missing or spurious values. Moreover, the updating and downdating procedures, based on solving inverse eigenvalue problems and eigenvalue problems, can be combined to yield a sliding window scheme, in which one node is replaced by another.

References

- [1] G.S. Ammar, W.B. Gragg and L. Reichel, On the eigenproblem for orthogonal matrices. In: *Proceedings of the 25th Conference on Decision and Control*, IEEE, New York, pp 1063–1066, 1986.
- [2] G.S. Ammar, W.B. Gragg and L. Reichel, Determination of Pisarenko frequency estimates as eigenvalues of an orthogonal matrix. In: F.T. Luk (Ed.), *Advanced Algorithms and Architectures for Signal Processing II*, Proc. SPIE 826, pp 143–145, 1987.
- [3] G.S. Ammar, W.B. Gragg and L. Reichel, Constructing a unitary Hessenberg matrix from spectral data. In: G.H. Golub and P. Van Dooren (Eds.), *Numerical Linear Algebra, Digital Signal Processing and Parallel Algorithms*, Springer, New York, pp 385–396, 1991.

- [4] G.S. Ammar, W.B. Gragg and L. Reichel, Downdating of Szegő polynomials and data fitting applications. *Lin. Alg. Appl.* **172**, pp 315–336, 1992.
- [5] G.S. Ammar, L. Reichel, and D.C. Sorensen, An implementation of a divide and conquer algorithm for the unitary eigenproblem. *ACM Trans. Math. Software*, to appear.
- [6] A. Bunse-Gerstner and L. Elsner, Schur parameter pencils for the solution of the unitary eigenproblem. *Lin. Alg. Appl.* **154–156**, pp 741–778, 1991.
- [7] A. Bunse-Gerstner and C. He, A Sturm sequence of polynomials for unitary Hessenberg matrices, preprint.
- [8] W.B. Gragg, Positive definite Toeplitz matrices, the Arnoldi process for isometric operators, and Gaussian quadrature on the unit circle (in Russian). In: E.S. Nikolaev (Ed.), *Numerical Methods in Linear Algebra*, Moscow University Press, Moscow, pp 16-32, 1982.
- [9] W.B. Gragg, The QR algorithm for unitary Hessenberg matrices. *J. Comput. Appl. Math.* **16**, pp 1–8, 1986.
- [10] W.B. Gragg and L. Reichel, A divide and conquer method for unitary and orthogonal eigenproblems. *Numer. Math.* **57**, pp 695–718, 1990.
- [11] L. Reichel and G.S. Ammar, Fast approximation of dominant harmonics by solving an orthogonal eigenvalue problem. In: J. McWhirter et al. (Eds.), *Proc. Second IMA Conference on Mathematics in Signal Processing*, Oxford University Press, pp 575–591, 1990.
- [12] L. Reichel, G.S. Ammar and W.B. Gragg, Discrete least squares approximation by trigonometric polynomials. *Math. Comp.* **57**, pp 273-289, 1991.
- [13] T.-L. Wang, *Convergence of the QR Algorithm with Origin Shifts for Real Symmetric Tridiagonal and Unitary Hessenberg Matrices*. Ph.D. Dissertation, Dept. of Mathematics, University of Kentucky, Lexington, KY, 1988.